

IVERMECTIN- ivermectin paste
JEFFERS

JEFFERS Ivermectin (ivermectin) Paste 1.87%

Anthelmintic and Boticide

Removes worms and bots with a single dose.

Contents will treat up to 1250 lb body weight.

FOR ORAL USE IN HORSES ONLY.

KEEP OUT OF REACH OF CHILDREN.

ANADA 200-326, Approved by FDA

Net Wt. 0.21 oz (6.08 g)

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. **JEFFERS®** Ivermectin Paste provides effective treatment and control of the following parasites in horses. **Large Strongyles**(adults) - *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; **Small Strongyles**(adults, including those resistant to some benzimidazole class compounds)- *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus* and *Petrovinema poculatum*; **Small Strongyles**- Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae)-*Oxyuris equi*; **Ascarids**(adults, third-stage and fourth-stage larvae)-*Parascaris equorum*; **Hairworms** (adults)-*Trichostrongylus axei*; **Large-mouth Stomach Worms**(adults)-*Habronema muscae*; **Bots**(oral and gastric stages)-*Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms**(adults and fourth-stage larvae)-*Dictyocaulus arnfeldi*; **Intestinal Threadworms**(adults)-*Strongyloids westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by **neck threadworm** microfilariae, *Onchocerca* sp.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. **JEFFERS®** Ivermectin Paste effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - **JEFFERS®** Ivermectin Paste kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. **JEFFERS®** Ivermectin Paste is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

☐**ANIMAL SAFETY:** **JEFFERS®** Ivermectin Paste may be used in horses of all ages, including mares

at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

STORE BETWEEN 20° C - 25° C (68° F - 77° F). Excursions between 15° C - 30° C (59° F - 86° F) are permitted.

WARNING: So not use in horses intended for human consumption.

Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain MSDS, contact JEFFERS at 1-800-JEFFERS.

PRECAUTIONS: JEFFERS® Ivermectin Paste has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with JEFFERS® Ivermectin Paste have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. microfilariae). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable.

Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with JEFFERS® Ivermectin Paste. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

JEFFERS® Ivermectin
(ivermectin) Paste 1.87%
Removes worms and bots with a single dose.
Contents will treat up to 1250 lb body weight.
FOR ORAL USE IN HORSES ONLY.
KEEP OUT OF REACH OF CHILDREN.
ANADA 200-326, Approved by FDA.
Net Wt. 0.21 oz (6.08 g)

Anthelmintic and Boticide

Manufactured for:
JEFFERS®
P.O. Box 100
Dothan, Alabama 36302

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. JEFFERS® Ivermectin Paste provides effective treatment and control of the following parasites in horses. **Large Strongyles** (adults)- *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicinctus* and *T. serratus* and *Craterostomum acuticaudatum*, **Small Strongyles** (adults), including those resistant to some benzimidazole class compounds)- *Coronocylus* spp. including *C. coronatus*, *C. labratus* and *C. ciliatus*, *Cyathostomum* spp. including *C. catinum* and *C. pateratum*, *Cylicocyclus* spp. including *C. ussini*, *C. leptostomum*, *C. nassarius*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. callitarsis*, *C. galei*, *C. longivittatus*, and *C. minutus* and *Petrovoina pectinata*; **Small Strongyles** - Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae)-*Oxyuris equi*; **Ascarids** (adults, third-stage and fourth-stage larvae)-*Parascaris equorum*; **Hairworms** (adults)-*Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults)-*Habronema muscae*; **Bots** (oral and gastric stages)- *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and

fourth-stage larvae)- *Dicyocaulus immitis*; **Intestinal Threadworms** (adults)-*Strongylidius westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp.; cutaneous third-stage larvae; Dermatitis caused by neck **threadworm** microfilariae, *Onchocerca* sp.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. JEFFERS® Ivermectin Paste effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - JEFFERS® Ivermectin Paste kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. JEFFERS® Ivermectin Paste is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: JEFFERS® Ivermectin Paste may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

STORE BETWEEN 20° C - 25° C (68° F - 77° F)
Excursions between 15° C - 30° C (59° F - 86° F) are permitted.

Manufactured by:
JEFFERS Animal Health Inc.
Cambridge, ON Canada N3B 2W4
Manufactured for:
JEFFERS
P.O. Box 100
Dothan, Alabama 36302
www.jeffers.com

REV. 10/13
PROOF OF PURCHASE
1040126100
1040126100

WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain MSDS, contact JEFFERS at 1-800-JEFFERS.

PRECAUTIONS: JEFFERS® Ivermectin Paste has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with JEFFERS® Ivermectin Paste have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. microfilariae). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable.

Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with JEFFERS® Ivermectin Paste. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

IVERMECTIN

ivermectin paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86010-100
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Ivermectin (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	Ivermectin	18.7 mg in 1 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86010-100-01	6.08 g in 1 CARTON		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200326	12/12/2013	

Labeler - JEFFERS (094385218)**Registrant** - Bimeda, Inc. (060492923)**Establishment**

Name	Address	ID/FEI	Business Operations
Bimeda-MTC Animal Health		256232216	manufacture

Revised: 12/2018

JEFFERS